

Yukon Immunization Schedule

V A C C I N E											
A G E	DTaP-HB-IPV-Hib (Diphtheria, Tetanus, Acellular Pertussis, hepatitis B, Inactivated Polio Vaccine, Haemophilus Influenza type B)	DTaP-IPV-Hib (Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine, Haemophilus Influenza type B)	DTaP-IPV / Tdap-IPV (Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine)	Rotavirus	Pneumococcal Conjugate	Meningococcal Conjugate	MMR (Measles, Mumps Rubella)	Varicella (Chicken pox)	HPV (Girls only)	Tdap (Tetanus, Diphtheria, Acellular Pertussis)	Influenza
2 months	✓			✓	✓	✓					
4 months	✓			✓	✓	‡					
6 months	✓				‡						✓*
12 months					✓	✓	✓	✓			
18 months		✓									
4-6 years			✓				✓	✓			
Grade 6	Usually administered in school.										
Grade 9	Usually administered in school.										
						✓***			✓**	✓	

‡ Children with specific medical conditions that place them at high risk of disease should receive an additional dose.

* Ages six months to nine years: a second dose required four weeks after first if receiving vaccine for first time.

** Starting 2014/2015 school year two doses; second dose six months after first dose.

*** Meningococcal Quadrivalent vaccine

Common side effects of immunizations include soreness and redness at the site of the injection, irritability/fussiness, and/or mild fever.

If your child experiences anything more severe, please contact your physician and report all severe reactions to one of the nurses at your local Health Centre.

For more information contact your community health centre to speak with a nurse.

August 2016